

C4 - Chapter 6 - Integration - Extra practice 2 - Solutions

$$1. a) \quad \frac{dy}{dx} = \frac{2xy}{1+x^2}$$

$$\int \frac{1}{y} dy = \int \frac{2x}{1+x^2} dx \quad \text{Let } u = 1+x^2$$

$$du = 2x dx$$

$$\Rightarrow \int \frac{1}{y} dy = \int \frac{1}{u} du$$

$$\ln|y| = \ln|u| + c$$

$$\ln|y| = \ln|1+x^2| + c$$

$$b) \text{ When } x=0, y=2 \Rightarrow \ln 2 = \ln 1 + c \Rightarrow c = \ln 2$$

$$\ln|y| = \ln|1+x^2| + \ln 2$$

$$\ln|y| = \ln|2(1+x^2)|$$

$$y = 2(1+x^2)$$

$$2. a) \quad x=t^2 \quad y=t^3$$

$$\frac{dx}{dt} = 2t \quad \frac{dy}{dt} = 3t^2 \quad \Rightarrow \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{3t^2}{2t} = \frac{3t}{2}$$

$$\text{When } t=1 \quad x=1, y=1, \frac{dy}{dx} = \frac{3}{2} \Rightarrow m_{\text{NORMAL}} = -\frac{2}{3}$$

$$\therefore y-1 = -\frac{2}{3}(x-1)$$

$$b) \text{ Area} = \int_0^1 y dx + \text{Area of triangle}$$

Now, normal will cut the x-axis at $x=5/2$ (found by setting $y=0$)

\therefore Triangle has height 1 and base $5/2 - 1 = 3/2$

$$\Rightarrow \text{Area} = \int_0^1 t^3 \cdot 2t dt + \frac{1 \times 3/2}{2} = \int_0^1 2t^4 dt + \frac{3}{4} = \left[\frac{2t^5}{5} \right]_0^1 + \frac{3}{4}$$

$$= \frac{2}{5} + \frac{3}{4} = \frac{23}{20}$$

$$3. \quad \int_2^4 x \ln x dx = \left[\frac{x^2}{2} \ln x \right]_2^4 - \int_2^4 \frac{1}{x} \cdot \frac{x^2}{2} dx = \left(\frac{4^2}{2} \ln 4 - \frac{2^2}{2} \ln 2 \right) - \int_2^4 \frac{x}{2} dx$$

$$u = \ln x \quad \frac{dv}{dx} = x$$

$$\frac{du}{dx} = \frac{1}{x} \quad v = \frac{x^2}{2}$$

$$= 8 \ln 4 - 2 \ln 2 - \left[\frac{x^2}{4} \right]_2^4$$


$$= 8 \ln 4 - \ln 2^2 - \left(\frac{4^2}{4} - \frac{2^2}{4} \right)$$

$$= 8 \ln 4 - \ln 4 - 3 = 7 \ln 4 - 3 \text{ AS REQUIRED}$$

4. a) $V = x^2 + 4x$

$$\frac{dV}{dt} = x^2 - 25$$

$$\frac{dV}{dx} = 2x + 4$$



$$\frac{dx}{dt} = \frac{dx}{dV} \cdot \frac{dV}{dt} = \frac{1}{2x+4} \cdot (x^2 - 25) = \frac{x^2 - 25}{2x+4} \quad \text{AS REQUIRED}$$

b) $\int \frac{2x+4}{x^2-25} dx = \int dt$

$$\frac{2x+4}{x^2-25} = \frac{2x+4}{(x-5)(x+5)} = \frac{A}{x-5} + \frac{B}{x+5}$$

$$2x+4 = A(x+5) + B(x-5)$$

$$x=5 \Rightarrow 14 = 10A \Rightarrow A = \frac{7}{5}$$

$$x=-5 \Rightarrow -6 = -10B \Rightarrow B = \frac{3}{5}$$

$$\int \frac{7/5}{x-5} + \frac{3/5}{x+5} dx = \int dt$$

$$\frac{7}{5} \ln|x-5| + \frac{3}{5} \ln|x+5| = t + C$$

When $t=0, x=0$ (since bottle was empty, hence height was 0)

$$\Rightarrow \frac{7}{5} \ln|-5| + \frac{3}{5} \ln 5 = 0 + C$$

$$C = 2 \ln 5$$

$$\Rightarrow \frac{7}{5} \ln|x-5| + \frac{3}{5} \ln|x+5| = t + 2 \ln 5$$

$$\Rightarrow t = \frac{7}{5} \ln|x-5| + \frac{3}{5} \ln|x+5| - 2 \ln 5$$

5a) i) $(\cos \theta + \sin \theta)^2 = \cos^2 \theta + 2 \cos \theta \sin \theta + \sin^2 \theta$
 $= 1 + \sin 2\theta$ (since $\sin^2 \theta + \cos^2 \theta = 1$)

ii) $\int_{\pi/4}^{\pi/2} (\cos \theta + \sin \theta)^2 d\theta = \int_{\pi/4}^{\pi/2} (1 + \sin 2\theta) d\theta = \left[\theta - \frac{1}{2} \cos 2\theta \right]_{\pi/4}^{\pi/2}$

$$= \left(\frac{\pi}{2} - \frac{1}{2} \cos \pi \right) - \left(\frac{\pi}{4} - \frac{1}{2} \cos \left(\frac{\pi}{2} \right) \right)$$

$$= \frac{\pi}{4} + \frac{1}{2} \quad \text{AS REQUIRED}$$

b) $\int_0^1 (2x+1)^4 dx = \left[\frac{(2x+1)^5}{5 \cdot 2} \right]_0^1 = \frac{3^5}{10} - \frac{1^5}{10} = 24.2$

$$6. \frac{2}{x(x+1)(x+2)} = \frac{A}{x} + \frac{B}{x+1} + \frac{C}{x+2} \quad \Rightarrow \quad 2 = A(x+1)(x+2) + Bx(x+2) + Cx(x+1)$$

$$x=0 \Rightarrow 2 = 2A \Rightarrow A=1$$

$$x=-1 \Rightarrow 2 = -B \Rightarrow B=-2$$

$$x=-2 \Rightarrow 2 = 2C \Rightarrow C=1$$

$$\therefore \frac{2}{x(x+1)(x+2)} = \frac{1}{x} - \frac{2}{x+1} + \frac{1}{x+2}$$

$$\int_2^4 \frac{2}{x(x+1)(x+2)} dx = \int_2^4 \left(\frac{1}{x} - \frac{2}{x+1} + \frac{1}{x+2} \right) dx = \left[\ln x - 2 \ln x+1 + \ln x+2 \right]_2^4$$

$$= \left[\ln x - \ln(x+1)^2 + \ln x+2 \right]_2^4 = \left[\ln \frac{x(x+2)}{(x+1)^2} \right]_2^4$$

$$= \ln \left(\frac{4(4+2)}{(4+1)^2} \right) - \ln \left(\frac{2(2+2)}{(2+1)^2} \right)$$

$$= \ln \left(\frac{24}{25} \right) - \ln \left(\frac{8}{9} \right) = \ln \left(\frac{24/25}{8/9} \right) = \ln \frac{27}{25}$$

$$= \ln 27 - \ln 25 = \ln 3^3 - \ln 5^2 = 3 \ln 3 - 2 \ln 5 \quad \text{AS REQUIRED}$$

$$7 a) \quad 2y = x - \sin x \cos x$$

$$2 \frac{dy}{dx} = 1 - (\cos^2 x - \sin^2 x) \quad \text{Implicit differentiation and product rule}$$

$$2 \frac{dy}{dx} = 1 - \cos^2 x + \sin^2 x$$

$$2 \frac{dy}{dx} = \sin^2 x + \sin^2 x \quad \Rightarrow \quad \frac{dy}{dx} = \sin^2 x \quad \text{AS REQUIRED}$$

$$b) \int \sin^2 x dx = \frac{x - \sin x \cos x}{2} + C$$

OK, b) is difficult. From a) we concluded that if $2y = x - \sin x \cos x$ (*)

$$\text{then } \frac{dy}{dx} = \sin^2 x \quad \Rightarrow \int dy = \int \sin^2 x dx$$

$$\Rightarrow y = \int \sin^2 x dx = \frac{x - \sin x \cos x}{2} + C \quad \text{using (*)}$$

$$c) \int x \sin^2 x dx = x \left(\frac{x - \sin x \cos x}{2} \right) - \int \frac{x - \sin x \cos x}{2} dx = x \left(\frac{x - \sin x \cos x}{2} \right) - \int \frac{x}{2} - \frac{1}{4} \sin 2x dx$$

$$u = x \quad \frac{dv}{dx} = \sin^2 x$$

$$= x \left(\frac{x - \sin x \cos x}{2} \right) - \left(\frac{x^2}{4} + \frac{1}{8} \cos 2x \right) + C$$

$$\frac{du}{dx} = 1 \quad v = \frac{x - \sin x \cos x}{2}$$

$$8. a) \quad \frac{dx}{dt} = -kx \quad \Rightarrow \quad \int \frac{1}{x} dx = \int -k dt$$

$$\ln|x| = -kt + C$$

$$\text{When } t=0, x=A \Rightarrow C = \ln A$$

$$\ln|x| = -kt + \ln A$$

$$x = e^{-kt + \ln A} = e^{-kt} \cdot e^{\ln A}$$

$$\Rightarrow x = Ae^{-kt} \quad \text{AS REQUIRED}$$

$$b) \quad \text{When } t=10, x = \frac{1}{3}A \Rightarrow \frac{1}{3}A = Ae^{-10k}$$

$$\frac{1}{3} = e^{-10k}$$

$$\ln\left(\frac{1}{3}\right) = -10k$$

$$k = -\frac{1}{10} \ln\left(\frac{1}{3}\right) = \frac{1}{10} \ln 3$$

$$\text{When } x = \frac{1}{2}A, \quad \frac{1}{2}A = Ae^{-t \cdot \frac{1}{10} \ln 3}$$

$$\frac{1}{2} = e^{-t \cdot \frac{1}{10} \ln 3}$$

$$\ln\left(\frac{1}{2}\right) = -t \cdot \frac{1}{10} \ln 3$$

$$t = \frac{-\ln\left(\frac{1}{2}\right)}{\frac{1}{10} \ln 3} = 6.31 \quad (\text{to 2 dp's})$$

$$9. a) \quad \int \sin 2y dy = -\frac{1}{2} \cos 2y + C$$

$$b) \quad \int xe^x dx = xe^x - \int e^x dx = xe^x - e^x + C$$

$$u = x \quad \frac{dv}{dx} = e^x$$

$$\frac{du}{dx} = 1 \quad v = e^x$$

$$c) \quad \frac{dy}{dx} = \frac{xe^x}{\sin y \cos y}$$

$$\int \sin y \cos y dy = \int xe^x dx$$

$$\int \frac{1}{2} \sin 2y dy = \int xe^x dx$$

$$-\frac{1}{4} \cos 2y = xe^x - e^x + C$$

$$10. a) \frac{5x^2 - 8x + 1}{2x(x-1)^2} = \frac{A}{x} + \frac{B}{x-1} + \frac{C}{(x-1)^2}$$

Careful!

$$5x^2 - 8x + 1 = 2A(x-1)^2 + 2Bx(x-1) + 2Cx$$

$$x=0 \Rightarrow 1 = 2A \Rightarrow A = \frac{1}{2}$$

$$x=1 \Rightarrow -2 = 2C \Rightarrow C = -1$$

$$x=-1 \Rightarrow 14 = 8A + 4B - 2C$$

$$14 = 4 + 4B + 2$$

$$8 = 4B \Rightarrow B = 2$$

$$\therefore \frac{5x^2 - 8x + 1}{2x(x-1)^2} = \frac{1/2}{x} + \frac{2}{x-1} - \frac{1}{(x-1)^2}$$

$$b) \int f(x) dx = \int \frac{1/2}{x} + \frac{2}{x-1} - \frac{1}{(x-1)^2} dx = \int \frac{1/2}{x} + \frac{2}{x-1} - (x-1)^{-2} dx$$

$$= \frac{1}{2} \ln|x| + 2 \ln|x-1| - \frac{(x-1)^{-1}}{-1} + C$$

$$c) \int_4^9 f(x) dx = \left[\frac{1}{2} \ln x + 2 \ln|x-1| + \frac{1}{x-1} \right]_4^9$$

$$= \left(\frac{1}{2} \ln 9 + 2 \ln 8 + \frac{1}{8} \right) - \left(\frac{1}{2} \ln 4 + 2 \ln 3 + \frac{1}{3} \right)$$

$$= \ln 3 + 6 \ln 2 + \frac{1}{8} - \ln 2 - 2 \ln 3 - \frac{1}{3} = 5 \ln 2 - \ln 3 - \frac{5}{24}$$

$$= \ln 32 - \ln 3 - \frac{5}{24} = \ln\left(\frac{32}{3}\right) - \frac{5}{24} \text{ AS REQUIRED}$$

$$11. a) \int x \sin 2x dx = -\frac{1}{2} x \cos 2x - \int -\frac{1}{2} \cos 2x dx = -\frac{1}{2} x \cos 2x + \int \frac{1}{2} \cos 2x dx$$

$$u = x \quad \frac{dv}{dx} = \sin 2x$$

$$\frac{du}{dx} = 1 \quad v = -\frac{1}{2} \cos 2x$$

$$= -\frac{1}{2} x \cos 2x + \frac{1}{4} \sin 2x + C$$

$$b) \frac{dy}{dx} = x \sin 2x \cos^2 y$$

$$\int \frac{1}{\cos^2 y} dy = \int x \sin 2x dx$$

$$\int \sec^2 y dy = \int x \sin 2x dx$$

$$\tan y = \frac{1}{2} x \cos 2x + \frac{1}{4} \sin 2x + C$$

$$\text{When } x = \pi/4, y = 0 \Rightarrow \tan 0 = \frac{1}{2} \frac{\pi}{4} \cos\left(\frac{\pi}{2}\right) + \frac{1}{4} \sin\left(\frac{\pi}{2}\right) + C \Rightarrow 0 = \frac{1}{4} + C \Rightarrow C = -\frac{1}{4}$$

$$\tan y = \frac{1}{2} x \cos 2x + \frac{1}{4} \sin 2x - \frac{1}{4}$$

$$12 \text{ a) } \int_1^2 \frac{2x^2+1}{x} dx = \int_1^2 \frac{2x^2}{x} + \frac{1}{x} dx = \int_1^2 2x + \frac{1}{x} dx = \left[\frac{2x^2}{2} + \ln x \right]_1^2$$

$$= (2^2 + \ln 2) - (1^2 + \ln 1) = 3 + \ln 2$$

$$\text{b) i) } \int x e^x dx = x e^x - \int e^x dx = x e^x - e^x + C = e^x(x-1) + C \quad \text{AS REQUIRED}$$

$$u = x \quad \frac{dv}{dx} = e^x$$

$$\frac{du}{dx} = 1 \quad v = e^x$$

$$\text{ii) Volume of revolution} = \pi \int_0^1 y^2 dx = \pi \int_0^1 (\sqrt{x} e^{x/2})^2 dx = \pi \int_0^1 x e^x dx$$

$$= \pi \left[e^x(x-1) \right]_0^1 = \pi \left(e^1(1-1) - (e^0(0-1)) \right)$$

$$= \pi$$

$$13 \text{ a) } \frac{8x+1}{(2x-1)(x+2)} = \frac{A}{2x-1} + \frac{B}{x+2}$$

$$8x+1 = A(x+2) + B(2x-1)$$

$$x=-2 \Rightarrow -15 = -5B \Rightarrow B=3$$

$$x=1/2 \Rightarrow 5 = \frac{5}{2}A \Rightarrow A=2$$

$$\therefore \frac{8x+1}{(2x-1)(x+2)} = \frac{2}{2x-1} + \frac{3}{x+2}$$

$$\text{b) } \frac{dy}{dx} = \frac{(8x+1)y}{(2x-1)(x+2)} \Rightarrow \int \frac{1}{y} dy = \int \frac{8x+1}{(2x-1)(x+2)} dx$$

$$\int \frac{1}{y} dy = \int \frac{2}{2x-1} + \frac{3}{x+2} dx$$

$$\ln|y| = \frac{2}{2} \ln|2x-1| + 3 \ln|x+2| + C$$

$$\text{When } x=1, y=1 \Rightarrow \ln 1 = \ln 1 + 3 \ln 3 + C \Rightarrow C = -3 \ln 3$$

$$\therefore \ln|y| = \frac{2}{2} \ln|2x-1| + 3 \ln|x+2| - 3 \ln 3$$

$$\ln y = \ln(2x-1) + \ln(x+2)^3 - \ln 27$$

$$\ln y = \ln \left(\frac{(2x-1)(x+2)^3}{27} \right)$$

$$y = \frac{(2x-1)(x+2)^3}{27}$$

14 a) $x=t+2$ $y=t^2+1$
 $dx=dt$

When $x=0$, $0=t+2 \Rightarrow t=-2$

When $x=2$, $2=t+2 \Rightarrow t=0$

$$V = \pi \int_{x=0}^{x=2} y^2 dx = \pi \int_{t=-2}^{t=0} (t^2+1)^2 dt \text{ AS REQUIRED}$$

b) $V = \pi \int_{-2}^0 t^4 + 2t^2 + 1 dt = \pi \left[\frac{t^5}{5} + \frac{2t^3}{3} + t \right]_{-2}^0 = \pi \left\{ 0 - \left(\frac{(-2)^5}{5} + \frac{2(-2)^3}{3} + (-2) \right) \right\} = \frac{206\pi}{15}$

15 a) i)

x	1	2	3
$y=12/x$	12	6	4

 $\int_1^3 \frac{12}{x} dx \approx \frac{1}{2} \cdot 1 \{ 12 + 2(6) + 4 \} = 14$

ii)

x	1	1.5	2	2.5	3
$y=12/x$	12	8	6	$24/5$	4

 $\int_1^3 \frac{12}{x} dx \approx \frac{1}{2} \cdot 0.5 \{ 12 + 2(8+6+24/5) + 4 \} = \frac{67}{5}$

b) $\int_1^3 \frac{12}{x} dx = [12 \ln x]_1^3 = 12 \ln 3 - 12 \ln 1 = 12 \ln 3$

i) % error = $\frac{|14 - 12 \ln 3|}{12 \ln 3} \times 100\% = 6.2\%$

ii) % error = $\frac{|\frac{67}{5} - 12 \ln 3|}{12 \ln 3} \times 100\% = 1.6\%$